

FULBRIGHT & JAWORSKI L.L.P.

A REGISTERED LIMITED LIABILITY PARTNERSHIP
666 FIFTH AVENUE, 31ST FLOOR
NEW YORK, NEW YORK 10103-3198
WWW.FULBRIGHT.COM

FACSIMILE TRANSMISSION

#27

DATE: August 7, 2003**MATTER NUMBER:** 10016357

NAME:	FAX No.:	PHONE No.:
Examiner Tam Davis Examiner Tony Caputa USPTO	1703 746-7145	1703 308-3995

FROM: Norman D. Hanson**PHONE:** (212) 318-3168**USER ID:** NH01030**FLOOR:** 24**RE:** LUB53537**NUMBER OF PAGES WITH COVER PAGE:** 5**Message:**

For 3:00 Interview.

CAUTION - CONFIDENTIAL

THE INFORMATION CONTAINED IN THIS FACSIMILE IS CONFIDENTIAL AND MAY ALSO CONTAIN PRIVILEGED ATTORNEY-CLIENT INFORMATION OR WORK PRODUCT. THE INFORMATION IS INTENDED ONLY FOR THE USE OF THE INDIVIDUAL OR ENTITY TO WHOM IT IS ADDRESSED. IF YOU ARE NOT THE INTENDED RECIPIENT, OR THE EMPLOYEE OR AGENT RESPONSIBLE TO DELIVER IT TO THE INTENDED RECIPIENT, YOU ARE HEREBY NOTIFIED THAT ANY USE, DISSEMINATION, DISTRIBUTION OR COPYING OF THIS COMMUNICATION IS STRICTLY PROHIBITED. IF YOU HAVE RECEIVED THE FACSIMILE IN ERROR, PLEASE IMMEDIATELY NOTIFY US BY TELEPHONE, AND RETURN THE ORIGINAL MESSAGE TO US AT THE ADDRESS ABOVE VIA THE U.S. POSTAL SERVICE. THANK YOU.

**IF YOU DO NOT RECEIVE ALL OF THE PAGES, PLEASE CALL
FANI MALIKOUZAKIS AT (212) 318-3220 AS SOON AS POSSIBLE.**

USPTO PATENT FULL-TEXT AND IMAGE DATABASE

Home	Quick	Advanced	Pat Num	Help
Hit List	Next	Bottom		
View Cart	Add to Cart			
Images				

(1 of 10)

United States Patent

6,488,932

Boon , et al.

December 3, 2002

Method for treating cancer by administering MAGE-derived peptides

Abstract

The invention involves the reception of particular nonapeptides by HLA molecules. The nonapeptides are derived from expression products of the MAGE gene family. The resulting complexes are identified by cytolytic T cells. Such recognition may be used in diagnostics, or therapeutically.

Inventors: Boon; Thierry (Brussels, BE); van der Bruggen; Pierre (Brussels, BE); De Plaen; Etienne (Brussels, BE); Lurquin; Christophe (Brussels, BE); Traversari; Catia (Milan, IT)

Assignee: Ludwig Institute for Cancer Research (New York, NY)

Appl. No.: 571263

Filed: December 12, 1995

Current U.S. Class: 424/185.1; 424/193.1; 424/277.1; 530/320; 530/328; 530/335; 530/402

Intern'l Class: C07K 005/00; A61K 039/395; A61K 039/00

Field of Search: 424/185.1,277.1,193.1 530/328,300,335,402

References Cited [Referenced By]

U.S. Patent Documents

<u>5342774</u>	Aug., 1994	Boon et al.	435/240.
<u>5405940</u>	Apr., 1995	Boon et al.	530/328.
<u>5462871</u>	Oct., 1995	Boon et al.	435/240.

Foreign Patent Documents

WO9202543	Feb., 1992	WO.
WO9220356	Nov., 1992	WO.
WO9403205	Feb., 1994	WO.

Other References

- Rothbard, J.B., Curr. Opin. Immunol. 2:99-105, Major histocompatibility complex-peptide interactions, 1989.*
- Engelhard, V. H., Curr. Opin. Immunol. 6:13-23, Structure of peptides associated with MHC Class I molecules, 1994.*
- Ochoa-Garay, J. et al., Mol. Immunol. 34(3):273-281, The ability of peptides to induce cytotoxic T cells in vitro does not strongly correlate with their affinity for the H-2L-d molecule: implications for vaccine design and immunotherapy, 1997.*
- Parham, P et al., Immunol. Rev. 143:142-180, The origins of HLA-A, B, C polymorphism, 1995.*
- Parker, K. C. et al., J. Immunol. 149:3580-3587, Sequence motifs important for peptide binding to the human MHC Class I molecule, HLA-A2, Dec. 1, 1992.*
- Salgaller, M. L. et al., Cancer Immunol. Immunother. 39:105-116, Generation of specific anti-melanoma reactivity by stimulation of human tumor-infiltrating lymphocytes with MAGE-1 synthetic peptide, 1994.*
- Smilek, D. et al., PNAS 88:9633-9637, A single amino acid change in a myelin basic protein peptide confers the capacity to prevent rather than induce experimental autoimmune encephalomyelitis, Nov. 1991.*
- Van der Zec, R. et al., Eur. J. Immunol. 19:43-47, Efficient mapping and characterization of a T cell epitope by the simultaneous synthesis of multiple peptides, 1989.*
- Rotzschke et al., "Isolated and analysis of naturally processed Viral peptides as recognized by cytolytic T cells", Nature 348:252-254 (1990).
- Van der Bruggen et al., "A Gene Encoding An Antigen Recognized By Cytolytic T Lymphocytes on a Human Melanoma", Science 254:1643-1647 (Dec. 1991).
- Traversari et al., "A Nonapeptide Encoded by Human Gene MAGE-1 Is Recognized on HLA-A1 By Cytolytic T Lymphocytes Directed Against Tumor Antigen MZ2-E". J.Exp.Med. 176: 1453-1457 (1992).

Primary Examiner: Chan; Christina Y.

Assistant Examiner: Lubet; Martha

Attorney, Agent or Firm: Fulbright & Jaworski, LLP

Parent Case Text

RELATED APPLICATIONS

This application is a divisional of Ser. No. 08/354,679, filed on Dec. 13, 1994, and pending, which is a divisional of Ser. No. 07/938,334, filed on Aug. 31, 1992, now U.S. Pat. No. 5,405,940.

Claims

We claimed:

1. A method for treating a patient afflicted with a tumor, neoplasm or cancer which expresses a tumor

rejection antigen precursor and presents HLA-A1 molecules on cell surfaces of said tumor, neoplasm or cancer, comprising administering to said patient an amount of a nonapeptide, the amino acid sequence of which is set forth at SEQ ID NO: 17, sufficient to form complexes between said nonapeptide and said HLA-A1 molecules and induce lysis of cells of said tumor, neoplasm, or cancer which present complexes of said nonapeptide and said HLA-A1 molecule by cytolytic T cells of said patient.

2. The method of claim 1, wherein said patient expresses tumor rejection antigen precursor MAGE-3.
3. The method of claim 1, wherein said nonapeptide is administered via direct injection, time release administration, or via coupling to a tumor specific antibody.
4. The method of claim 1, wherein said cancer is melanoma.

Description

FIELD OF THE INVENTION

This invention relates to immunogenetics and to peptide chemistry. More particularly, it relates to a class of nonapeptides useful in various ways, such as immunogens and as materials which target and bind MHC/HLA molecules, as well as a cellular model useful in the testing of peptides and other molecules as vaccines, especially cancer vaccines. Most particularly, it relates to the so-called "tumor rejection antigens", in particular the MAGE family of these antigens.

BACKGROUND AND PRIOR ART

The study of the recognition or lack of recognition of cancer cells by a host organism has proceeded in many different directions. Understanding of the field presumes some understanding of both basic immunology and oncology.

Early research on mouse tumors revealed that these displayed molecules which led to rejection of tumor cells when transplanted into syngeneic animals. These molecules are "recognized" by T-cells in the recipient animal, and provoke a cytolytic T-cell response with lysis of the transplanted cells. This evidence was first obtained with tumors induced in vitro by chemical carcinogens, such as methylcholanthrene. The antigens expressed by the tumors and which elicited the T-cell response were found to be different for each tumor. See Prehn, et al., J. Natl. Canc. Inst. 18: 769-778 (1957); Klein et al., Cancer Res. 20: 1561-1572 (1960); Gross, Cancer Res. 3: 326-333 (1943), Basombrio, Cancer Res. 30: 2458-2462 (1970) for general teachings on inducing tumors with chemical carcinogens and differences in cell surface antigens. This class of antigens has come to be known as "tumor specific transplantation antigens" or "TSTAs". Following the observation of the presentation of such antigens when induced by chemical carcinogens, similar results were obtained when tumors were induced in vitro via ultraviolet radiation. See Kripke, J. Natl. Canc. Inst. 53: 333-1336 (1974).

While T-cell mediated immune responses were observed for the types of tumor described supra, spontaneous tumors were thought to be generally non-immunogenic. These were therefore believed not to present antigens which provoked a response to the tumor in the tumor carrying subject. See Hewitt, et al., Brit. J. Cancer 33: 241-259 (1976).

The family of tum.sup.- antigen presenting cell lines are immunogenic variants obtained by mutagenesis of mouse tumor cells or cell lines, as described by Boon et al., J. Exp. Med. 152: 1184-1193 (1980), the

(D) TOPOLOGY: linear
(ii) MOLECULE TYPE: protein
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15
Glu Val Val Pro Ile Ser His Leu Tyr
5

(2) INFORMATION FOR SEQ ID NO: 16:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 9 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
(ii) MOLECULE TYPE: protein
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 16
Glu Val Val Arg Ile Gly His Leu Tyr
5

(2) INFORMATION FOR SEQ ID NO: 17:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 9 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
(ii) MOLECULE TYPE: protein
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 17
Glu Val Asp Pro Ile Gly His Leu Tyr
E V D P SI G H L Y

(2) INFORMATION FOR SEQ ID NO: 18:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 9 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
(ii) MOLECULE TYPE: protein
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 18
Glu Val Asp Pro Ala Ser Asn Thr Tyr
5

(2) INFORMATION FOR SEQ ID NO: 19:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 9 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
(ii) MOLECULE TYPE: protein
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 19
Glu Val Asp Pro Thr Ser Asn Thr Tyr
5

(2) INFORMATION FOR SEQ ID NO: 20:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 9 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
(ii) MOLECULE TYPE: protein
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 20
Glu Ala Asp Pro Thr Ser Asn Thr Tyr
5

(2) INFORMATION FOR SEQ ID NO: 21:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 9 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single